



E-ISSN: 2707-2835
P-ISSN: 2707-2827
IJPLS 2020; 1(2): 29-32
Received: 14-05-2020
Accepted: 18-06-2020

Hamida ALbarasi
M.B.Ch.B, MSc,
Pharmacology and lecturer
assistant at Department of
Pharmacology, faculty of
drugs, Benghazi University,
Benghazi, Libya

Samia Elzwi
M.B.Ch.B, MSc,
Pharmacology and Assistant
professor at Department of
Pharmacology, faculty of
drugs, Benghazi, Libya

Corresponding Author:
Hamida ALbarasi
M.B.Ch.B, MSc,
Pharmacology and lecturer
assistant at Department of
Pharmacology, faculty of
drugs, Benghazi University,
Benghazi, Libya

Effect of cinnamon plant extract on blood sugar in mice

Hamida AL barasi and Samia Elzwi

DOI: <https://doi.org/10.33545/27072827.2020.v1.i2a.15>

Abstract

Cinnamon may be a small evergreen tree 10-15 meters tall, belongs to the family Lauraceae. The leaves are ovate-oblong in shape, 7-18 Cm long, the flowers which are arranged in panicles, have greenish color and distinct odor. The fruit may be a purple one centiberry containing one seed. Cinnamon are widely used as spice and flavoring agent and may be clinically utilized in treatment of certain diseases like type 2 DM. Cinnamon extract has blood sugar lowering effect in patients with type 2 diabetes. This study tested the effect of cinnamon extract on D- glucose induced hyperglycemias which showed significant reduction in plasma glucose level. Chronic hyperglycemia model induced by one intraperitoneal injection of streptozotocin, the extract showed marked decrease in blood sugar level.

Keywords: cinnamon extract; diabetes mellitus; d- glucose; experimental animals; and streptozotocin

Introduction

DM is disease characterized by hyperglycemia it's related to relative or absolute impairment in insulin secretion, along side varying degree of resistance to the action of insulin.

The American Diabetes Association (ADA) issued diagnostic criteria for DM in 1997 with follow up in 2003. The diagnostic criteria are supported one among three following abnormalities: fasting plasma glucose, random elevated glucose with symptoms, or abnormal oral glucose tolerance test as shown in table 1 [1]. the 2003 criteria lowered the fasting plasma glucose level went to define impaired fasting glucose (IFG) from the 1997 criteria but maintained an equivalent definitions for diabetes and impaired glucose tolerance (IGT) as measured by an oral glucose tolerance test. Patients with IFG and / or IGT are now mentioned as having pre-diabetes indicating the relatively high risk for development of diabetes in these.

There are two sorts of diabetes type 1 is characterized by destruction of the pancreatic B cells resulting in absolute insulin deficiency. this is often usually thanks to autoimmune destruction of B cells (Type 1A). pancreatic

Testing for Islet- cell antibodies (ICA) or other autoantibodies (antibodies to glutamic acid dehydrogenase [ant-GAD] and anti-insulin) in serum could also be helpful if establishing the diagnosis is important: a positive result's indicative of type1A diabetes. However, some patients haven't any evidence of autoimmunity and haven't any other cause for cell destruction (idiopathic type 1B DM [2].

Type 2 diabetes is liable for over 80% of cases of diabetes within the us, Canada, and Europe. Type 1 diabetes accounts for an additional 5 to 10 %, with the rest thanks to other causes. New information has led to increased understanding of genetic defects associated with diabetes, and therefore the formerly termed MODY syndrome (maturity onset diabetes of the young) have now been linked to specific genetic defects [3].

Cinnamon has been known from remote antiquite, and it had been so highly prized among ancient nation that it had been considered a present for monarchs and other great potentates. it had been imported to Egypt from China as early as 2000BC [4]. within the middle Ages, the source of cinnamon was a mystery to the Western world Arab traders brought the spice via overland trade route to Alexandria in Egypt, where it had been bought by Venetian traders from Italy who held a monopoly on the spice trade Europe [5]. Its flavor is thanks to an aromatic volatile oil which makes up 0.5 to a quarter of its composition.

Table 1: Criteria for diagnosis of DM

1.	1-Symptoms of diabetes and an off-the-cuff plasma glucose 200 mg/dl (11.1 mmol/l). casual is defined as any time of day without reference to time since last meal. The classical symptoms of DM include polyuria, polydipsia, and unexplained weight loss
2.	2-FPG126 mg/dl (7.0 mmol/l). Fasting is defined as no caloric intake for a minimum of 8 hours
3.	- 3-2 hrs. plasma glucose 200mg/dl (11.1 mmol/l) during an oral glucose tolerance test. The test should be performed as described by WHO using glucose load containing the equivalent of 75 g anhydrous glucose dissolved in water.

This oil is ready by roughly pounding the bark, macerating it within the sea – water, then quickly distilling the entire [6]. it's of a golden yellowish color with characteristic odors of cinnamon and a really hot aromatic taste. The pungent taste and scent come from cinnamic aldehyde or cinnamaldehyde and by the absorption of oxygen because it ages, it darkens color and develops resinous compounds. Chemical composition of volatile oil includes ethyl cinnamate, eugenol, cinnamaldehyde, beta- caryophyllene, linalool and methyl chavicol [7]. Cinnamon leaf oil features a warm, spicy, but rather harsh odors, lacking the rich body of the bark oil [8].

Its major constituents are eugenol instead of cinnamaldehyde, Its used as flavoring agents for seasoning and savory snacks. As an inexpensive fragrance it's added to soaps. The oils high eugenol content also makes it valuable as source of this chemical subsequent conversion into iso-eugenol. another flavoring agents.

Glucose is that the most abundant monosaccharide, a subcategory of carbohydrates. The present sort of glucose is D-glucose, while L-glucose is produced synthetically in comparatively small amounts and is of lesser importance. Glucose may be a monosaccharide containing six carbon atoms and an aldehyde radical and is therefore mentioned as an aldohexose. (C₆H₁₂O₆) [9].

Streptozotocin (STZ, 2- deoxy-2- (3 methyl 3-nitrosoureido) D – glucopyranose) is synthesized by streptomycetes achromogenes and is employed to induce both insulin dependent and non-insulin dependent DM (IDDM and NIDDM respectively). The frequently used single intravenous dose in rat to induce IDDM is between 40- 60 mg/kg b.w [10], but higher dose also are used.

Materials and methods

Preparation of plant extract

Dried Cinnamon Bark was obtained from the local market, crushed by using electrical crusher to get the powder. The solvent extraction was performed by using petroleum ether, chloroform, ester, ethanol and water to urge the aqueous extract of the plant. This was done by using continuous percolation process and therefore the method used for this, was Soxhlet apparatus extraction [11].

500 grams of coarse plant powder were put within the thimble siphons of Soxhlet after maceration within the same solvent for 15 minute, the flask of Soxhlet was crammed with about one liter of the used solvent and heating of solvent was started at temperature 60-80 C°. And by condensation process the solvent within the thimble siphons off into main vessel containing the extract. The

method continued for 4-5 times until complete extraction was achieved. The solvent from the flask was evaporated by using the rotatory evaporator under vacuum at temperature of 40-50 °C and rotation was continued until the extract of specific solvent was obtained

The extraction process started by using solvent with less polarity and gradually increasing the polarity with other solvents as follow

Petroleum ether, chloroform, ester, ethanol and water. Finally, 20 grams of dry powder were obtained from each complete extraction cycle, the percent of yield was 4%

Effect of cinnamon extract on acute hyperglycemia induced by D- glucose

Male albino mice were used, animals were divided into seven groups each of six animals. The primary group was control group it had been treated with normal saline (0.2 ml/mouse I.P). The second group was given one dose of D – glucose (10 g/kg, IP) it had been the positive control group [12]. The third, fourth, fifth, sixth and seventh groups were treated with single dose of cinnamon extract (300mg/kg PO) at same time. Later on, these five groups were injected by D - glucose (10 g/kg IP) at different time intervals from the treatment with cinnamon extract at 1, 3, 6,10, 24 hrs.). The blood sugar level was estimated altogether animal groups by using Accu- cheek glucose test strip this was by piercing the tail of the mice, a drop of blood was placed on a blood sugar test strip which then inserted into glucose meter.

Estimation of the effect of cinnamon extract on chronic hyperglycemia induced by streptozotocin

To induce diabetes female mice were injected with streptozotocin (100 mg/kg IP) streptozotocin was dissolved in 0.1 mol / L sodium citrate buffer (pH4.0) and injected within 5 minute [13]. to work out the consequences of cinnamon extract mice were divided into five groups each of six animals the primary non-treated control group was given a citrate buffer (0.5 ml/mouse PO) and therefore the third group was treated only with cinnamon extract (300 mg/kg PO) for two weeks (positive control group). The second, fourth and fifth biological group were fasted about 12 hours prior treatment with single dose of streptozotocin (100 mg/kg IP).

The fourth group was pretreated with cinnamon extract (300mg/kg /day PO) for 2 weeks before treatment with streptozotocin and lastly the fifth group treated with one dose of glibenclamide (0.5 mg/kg PO) [14]. 2 hours before collecting blood on the day of sacrifice mice were kept overnight fasted then decapitated without anesthesia and trunk blood was collected into perhilled tubes containing 1 mg/ml of EDTA for insulin and glucose determination. Blood sugar decided using glucose oxidase kit, plasma insulin was assayed employing a standard radioimmunoassay technique.


Statistical Analysis: The results of this experiment was done by use of descriptive analysis because the mean S.E.M. Data was analyzed by using T- test in some experiment and therefore the method of study of variance (ANOVA). The many of study is indicated by the p value if it's adequate to but 0.05 it's considered as statistically significant.

Results

Hypoglycemic effect of cinnamon extract on D- glucose induced hyperglycemia

As shown in figure (1), the plasma glucose level at different time intervals from the administered single Cinnamon

extract dose was significantly decreased after 6 hours as compared to thee all other groups (control, zero time, at 1 hour, at 3 hour, at 10 hour ad at 24 hours) and therefore the p value was ($P < 0.001$).




** Singnificantly decreased as compared to all other groups ($P < 0.001$).

Fig 1: Effect of CE on acute hyperglycemia induced by D- Glucose Administered at Different time intervals

The effect of cinnamon extract on streptozotocin induced hyperglycemia

As shown in figure (2) by using ANOVA followed by post hoc ergo propter hoc test, our data indicated that, there was a big increase within the level of glucose within the streptozotocin treated groups as compared to all or any other groups. ($P < 0.001$).

On other hand the pretreated cinnamon extract group showed a big decrease within the glucose level as compared to the streptozotocin group (p, 0.001), also the glibenclamide significantly reduced the glucose level as compared to the streptozotocin group ($P < 0.005$).



** Singnificantly increased as compared to control groups ($P < 0.05$).

* Singnificantly decreased as compared to STZ groups ($P < 0.001$).

Fig 2: Effect of CE and Glibenclamide (GC) hyperglycemia induced by A single dose of STZ in Mice.

Discussion

Cinnamon has been widely used as spice and flavoring agent, some previous studies have been shown that it may be clinically useful in treatment of certain diseases such as type 2 diabetes mellitus, dyspepsia, anorexia, inflammatory disorders in addition to its antimicrobial activity [15].

The effect of cinnamon extract on hyperglycemia was investigated using 2 animal models namely D –glucose model) and chronic streptozotocin model. Previous studies suggested that D – glucose is mediated by the stimulation of the lateral part of the dorsal parabrachial nucleus (LPBD) enhancing glucagon release and suppression insulin

secretion leading to hyperglycemia. D – glucose affects the suprachiasmatic nucleus (SCN) and β –adrenergic mechanisms that are involved in the hyperglycemic response to (LPBD) stimulation [16].

Pretreatment with oral cinnamon extract had a weak and delayed hypoglycemic effect which was significant 6 hour after treatment. This transient and delayed effect indicated that the cinnamon extract is slowly absorbed and therefore it may have low bioavailability. In addition the hypoglycemic effect which occurred only after 6 hour may be explained through an extra pancreatic mechanism, this may imply that cinnamon extract may affect (LPBD or SCN) nuclei in hypothalamus and therefore counteracting the effect of D – glucose by suppressing glucagon and enhancing insulin secretion. More over methyl hydroxyl chalone (MHCP) one of the component of cinnamon extract was demonstrated in vivo to activate glycogen synthase and inhibit glycogen synthase kinase 3 B [17].

Streptozotocin may be given in multiple low doses for induction of IDDM in mouse such effect is mediated by activation of immune function. However [18] demonstrated that the nonspecific activation of immune system via complete Freud's adjuvant prior to streptozotocin injection allows to reduce its diabetogenic dose even in the rat.

NIDDM can be easily induced in rats by intravenous or intraperitoneal treatment with 100 mg/kg b.w.

Streptozotocin action in the B cell is accompanied by characteristic alteration in blood insulin and glucose concentration. Two hours after injection, the hyperglycemia is observed with a concomitant drop in blood insulin, about six hours later hyperglycemia occurs with high levels of blood insulin. Finally, hyperglycemia develops and blood insulin levels decrease. These changes in blood glucose and insulin concentrations reflects abnormalities in B cell function.

A study was carried out in Jordan on 75 patients with type 2 diabetes for 4 weeks; the patients were asked to take 2 g of ground cinnamon (two 500 milligrams of cinnamon capsules) immediately after breakfast, lunch, and dinner; and, as a result, a daily dose of 6 g of cinnamon was found to be effective in reducing blood glucose in a short time [19].

In a parallel study on patients with type 2 diabetes after menopause, the patients were given cinnamon (*Cinnamomum cassia*, 1.5 g/day) and placebo supplements for 6 weeks. Based on the study, no change was detected in preprandial blood glucose, preprandial insulin and HbA1c levels, blood lipids, and whole-body insulin resistance/sensitivity^[20]. In another study, 25 postmenopausal women were examined. The consumption of 1500 mg of cinnamon per day has been observed not to reduce blood sugar compared to placebo^[21].

CIN effects were investigated on fasting blood glucose, insulin, oral glucose tolerance test (OGTT), insulin tolerance test (ITT), liver biomarkers, lipid profile, oxidative stress biomarkers, serum advanced glycation end products (AGEs) and its receptors (RAGE) in the aorta, and histopathology of the liver and aorta. Additionally, the mRNA expression of hepatic insulin signaling pathway genes, phosphorylated AKT (serine 473) (P-AKT ser473) level, and aortic nitric oxide synthase3 (eNOS) and NADPH oxidase4 (NOX4) were determined. CIN treatment for 30 days significantly decreased OGTT, ITT, fasting blood glucose, insulin, and HOMA-IR and increased HOMA- β index when compared to diabetic rats. CIN also improved lipid profile and decreased serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activity, serum AGEs, and aortic RAGE. Additionally, CIN treatment significantly decreased hepatic malondialdehyde (MDA), increased hepatic and aortic glutathione (GSH) and superoxide dismutase (SOD), and decreased steatosis and inflammation observed in liver tissue of rats. Additionally, significant elevation in mRNA expression of insulin receptor substrate1 (IRS1), phosphatidylinositol 3-kinase regulatory subunit1 (PI3K-P85 subunit), and AKT serine/threonine kinase2 (AKT2); increased levels of P-AKT ser473 and aortic eNOS; and decrease in mRNA expression of NOX4 were detected in CIN-treated group when compared to diabetic group. This study suggests antidiabetic and antioxidant effects of CIN probably through upregulation of eNOS and IRS1/PI3K/AKT2 signaling pathway and alleviating AGEs, RAGE, and NOX4 elevation^[22].

References

- Genuth S, Albert KG, Bennett P. Follow up report on the diagnosis of diabetes mellitus. *Diabetes Care* 2003;26:3160.
- Liberman IM, Pietropaolo M, Trucco M. Islet cell autoimmune in white and black children and adolescents with IDDM. *Diabetes Care* 1998;21:1824.
- Rother KI. Diabetes treatment – Bridging the divide *N Engl J Med* 2007;356(15):1499-1501.
- Jayaprakasha GK, Jagan Mohan Rao L, Sakariah KK. Volatile constituents from cinnamon *zeylanicum* fruit stalks and their antioxidant activities. *J Agric Food Chem* 2003;51(15):4344-4348.
- GruenwaldJ. PDR for Herbal Medicine 3rd ed, Montvale NJ Thompson PDR; 199- 200.arvil- Tylor, K.J Anderson R.A, and Graves, D.J (2001). A hydroxychalcone deived from cinnamon function as amemetic for insulin in 3T3-L1 adipocytes. *J. Am. Coll. Nutr.* 2004;20:327-336.
- Moestafa A, Badeges F. Distillation of cassia leaf oils (*Cinnamon burmanii*, *C. Zeylanicum* and *C.cassia*) by cohobation method and identification of oil components. *Warta Industri Hadil Pertanian Indonesia* 1986;3(1):22-25.
- Arpi A, Anahit M, Margarit P, Naira S, Anush B et al Chemical composition and some biological activities of the essential oils from basil *Ocimum* different cultivars *BMC Complementary and Alternative Medicine* 2017;17:PMC5248495.
- Lockwood GB. Major constituents of the essential oils of Cinnamon cassia Blume growing in Nigeria. *Planta Medica* 1979;36(4):380-381.
- Purseglove JW, Brown EG, Green CL, Robbins SRJ. Cinnamon and cassia ; pp 100-173. In *spices* 1981;1:439. London: Longman.
- Ganda OP, Rossi AA, Like AA. Studies on streptozotocin diabetes. *Diabetes* 1976;25:595-603.
- Trease and Evans pharmacology. Comparison of Phytochemicals and Anti-Nutritional Factors in Some Selected Wild and Edible Bean in Nigeria. *Food and Nutrition Sciences* 1978;(7):11TH. ED
- ElDebani AH, El-Darrat A, Abdellatif AG. Reversal of drug induced hyperalgesia by transient hyperglycemia. *Proceedinh of the third Jamahiriya Conference of Medical Sciences. Tripoll*, 31 st May 3 rd June 1996.
- Beom K, Eun-Kyung K, Eun-Sil J, Do-Go R. Cortex cinnamon extract prevents streptozotocin- and cytokine – induced B- cell damage by inhibiting NF-Kb. *World J Gastroenterol* 2006;12(27):4331-4337.
- Jean-Jacques D, Dugald S, Dan P, Kieran C, Taryan F, Edward M, *et al.* From type 2 diabetes to antioxidant activity: A common and cassia cinnamon bark *Can.J Physiol. Pharmacol* 2007;85:837-847.
- Fujiwara, K. Nagai, S. Takagi S. and H. Nakagawa. Hyperglycemia induced by electrical stimulation of lateral part of the dorsal parabrachial nucleus *Am J Physiol Endocrinol Metab* 1988;254:E468-E475.
- Jarvil-Taylor KJ, Anderson RA, Graves DJ. A hydroxychalcone derived from cinnamon function as amimetic for insulin in 3T3-L1 adipocytes. *J. Am. Coll. Nutr* 2001;20:327-336.
- Ziegler M, Zeigler B, Hehmke B. Sever hyperglycemia caused by autoimmunization to B – cells in rats *Diabetologia* 1984;27:163-165.
- Portha B, Levacher C, Picon L, Rosselin G. Diabetogrnic effet of streptozotocin in the rat during the perinatal period. *Diabetes* 1974;23:889-895.
- Al Jamal AR. Effects of cinnamon on blood glucose and lipids levels in diabetic patients (type1),” *African Journal of Biochemistry Research* 2009;3:181-184. View at: Google Scholar
- Rudkowska. Functional foods for health: focus on diabetes,” *Maturitas* 2009;62(3):263-269. View at: Publisher Site | Google Scholar
- Vanschoonbeek K, Thomassen BJW, Senden JM, Wodzig WKWH, Van Loon LJC. Cinnamon supplementation does not improve glycemic control in postmenopausal type 2 diabetes patients,” *Journal of Nutrition* 2006;136(4):977-980. View at: Publisher Site Google Scholar
- Marwa EA, George S, Rami A, AGhada MS, Hatem A Salem. Cinnamaldehyde ameliorates STZ-induced rat diabetes through modulation of IRS1/PI3K/AKT2 pathway and AGEs/RAGE interaction *Naunyn-Schmiedeberg's Archives of Pharmacology* 2019;392:243-258. Cite this article.