



E-ISSN: 2707-2835

P-ISSN: 2707-2827

www.pharmacognosyjournal.com

IJPLS 2021; 2(1): 44-48

Received: 20-11-2020

Accepted: 24-12-2020

Gaurava Srivastav

Research Scholar, Department of Pharmacology, Advance Institute of Biotech and Paramedical Sciences Kanpur, Uttar Pradesh, India

Dakshina Gupta

Assistant Professor, Department of Pharmacology, Advance Institute of Biotech and Paramedical Sciences Kanpur, Uttar Pradesh, India

Corresponding Author:

Gaurava Srivastav

Research Scholar, Department of Pharmacology, Advance Institute of Biotech and Paramedical Sciences Kanpur, Uttar Pradesh, India

Review on cinnamon plant

Gaurava Srivastav and Dakshina Gupta

DOI: <https://doi.org/10.33545/27072827.2021.v2.i1a.26>

Abstract

The goal of this article is to use scholarly search engines like PubMed and Scifinder to find scholarly articles and reports on Cinnamon, its novel implications, preparation, analysis, and use in the management and cure of serious illnesses like diabetes, cardiovascular disease, Alzheimer's disease, and cancer. Cinnamon has long been used in food preparations and as a herbal medication to treat a wide range of diseases and symptoms. Cinnamon has antioxidant, antibacterial, anti-inflammatory, and other medicinal effects. New research not only demonstrates the value of cinnamon as a flavor, but also suggests that it could be used as a natural treatment for significant ailments like type 2 diabetes, chronic digestive issues, cardiovascular disease, cancer, and Alzheimer's disease. This article examines the botanical, chemical, and pharmacological properties of cinnamon in depth.

Keywords: cinnamon, pubmed, diabetes, cardiovascular disease, alzheimer's disease

Introduction

Cinnamon is made from the inner bark of tropical evergreen cinnamon trees, and its name comes from a Greek phrase that means "sweet wood."^[1] *Cinnamomum* (cinnamon) is a Lauraceae genus with a large number of members used as spices^[2]. The Ceylon or real cinnamon (*Cinnamomum zeylanicum* Blume), which is found in Sri Lanka and Southern India (Fig. 1A), and the cassia (*Cinnamomum aromaticum* Ness), which is grown in China, Indonesia, and Vietnam, are the two principal types of cinnamon (Fig. 1B).

When dried, cassia forms a tubular form called a quill or cinnamon stick. Since ancient times, Egyptians and Chinese have employed cinnamon in food preparations and traditional medicine.⁴ Furthermore, this spice has been discovered to have potent antioxidant, antibacterial, antipyretic^[5], and anti-inflammatory effects, all of which aid tissue repair^[6]. In oriental herbal medicine, cinnamon bark has been used as a spice and to produce tea, as well as a herbal therapy for treating common colds, cardiovascular diseases, and chronic gastrointestinal and gynaecological issues.⁷ Sore throats, coughs, indigestion, abdominal pains, in-testinal spasms, nausea, flatulence, and diarrhoea have all been treated with cinnamon. Moreover, it has been found that cinnamon slows down food spoilage and displays antifungal properties^[8]. One study has found that cinnamon bark water extract enhances glucose uptake by increasing insulin sensitivity. Several studies have also found that cinnamon extract (CE) displays anticancer activity^[5] and inhibits angiogenesis by blocking vascular endothelial growth factor (VEGF) 2 signaling. These results indicate that the observed pharmacological activities may have originated from the polyphenolic compounds in cinnamon water extract^[7].

Components

Cinnamon is a plant with many uses as an herbal medicine and it contains mucilage, tannin, sugar, resin, and essential oil, among which essential oil is the most important constituent. The major portion of the essential oil is made up of cinnamaldehyde or cinnamic aldehyde^[6]. Cinnamaldehyde is responsible for the flavor and aroma of cinnamon^[9]. A study found that cinnamaldehyde is the major volatile component of cinnamon sticks, with 83.6% of essential oils extracted from cinnamon stick powder^[2]. The essential oils from Cinnamon cassia also contain 80e90% cinnamaldehyde with little or no eugenol, which is different from *C. zeylanicum* bark, which contains 60e80% cinnamaldehyde and approximately 2% eugenol, however, essential oils from its leaves were found to be rich in eugenol (70e75%)^[2].

The study showed that the crude extracts of cinnamon stick also contained high levels of nonvolatile compounds (mainly condensed tannins), which consist of 23.2% proanthocyanidins and 3.6% catechins in addition to cinnamaldehyde (64.1%)^[12].




Fig 1: (A) *True-cinnamon*. (B) *Cassia cinnamon*

Diabetes

Diabetes mellitus is a chronic disorder of glucose metabolism caused by the dysfunction of pancreatic beta cells and insulin resistance. Diabetes is a serious health problem that affects both sexes in all age groups and all over the world^[10]. Cinnamon has been shown to have insulin mimetic properties because its biologically active substances enhance glucose uptake by activating insulin receptor kinase activity, autophosphorylation of the insulin receptor, and glycogen synthase activity^[11]. In one study, a factor called methyl hydroxyl chalcone polymer, which was isolated from cinnamon, was observed to increase insulin-dependent glucose metabolism approximately 20-fold *in vitro*. Safdar *et al.*^[10] explained that methyl hydroxyl chalcone polymer made fat cells more responsive to insulin by activating the enzyme that causes insulin to bind to cells and inhibiting the enzyme that blocks the process leading to maximal phosphorylation of the insulin receptor, which is associated with increased insulin sensitivity.

Cinnamon is rich in polyphenolic components, and it was shown that an intake of 500 mg/d for 12 weeks decreased oxidative stress and improved impaired fasting glucose^[12]. Another finding also indicated that the consumption of 500 mg/d of a specific aqueous extract of cinnamon (Cinnulin PF) for a 12-week period significantly improved some metabolic syndromes such as fasting blood sugar, systolic blood pressure, and body composition, and because there is a relationship between metabolic syndrome and cardiovascular disease, these results can have an important effect for the improvement of public health^[13].

Anticholesterol

Hyperlipidemia is described by raised serum levels of total cholesterol (TC), low-density lipoprotein (LDL), Very low-density lipoprotein (VLDL), and diminished serum level of high density-lipoprotein (HDL). According to American heart incorporation, a high level of fats known as hyperlipidemia. These fats consist of cholesterol and triglyceride. Lipids and fatty substances in the blood and is a greater risk factor in the growth of atherosclerosis and heart diseases^[29].

The compounds present in Cinnamon have shown to be beneficial for glucose uptake, insulin regulation, and bloodlipid profile. A study was performed on patients with type 2 diabetes consuming cinnamon. The results showed that there were significant decreases in fasting serum glucose (18e29%), tri-glycerides (23e30%), total cholesterol (12e26%), and low-density lipoprotein cholesterol (7e27%) for patients who consumed 1 g, 3 g, or 5 g of cinnamon for 40 days^[4, 14].

Antioxidant

Epicatechin, camphene, eugenol, gamma-terpinene, phenol, salicylic acid, and tannins, which are among the natural dietary antioxidants contained in cinnamon, have been demonstrated to lower oxidative stress in a dose-dependent manner via the inhibition of 5-lipoxygenase^[15]. Natural antioxidants are receiving a lot of attention, and efforts to replace synthetic antioxidants are ongoing.

Furthermore, these natural antioxidants can be packaged into functional meals, which can assist the body avoid oxidative damage^[4].

Cancer

Cancer remains to be a major global healthcare burden as 18.1 million new cases and 96 million death occurred 2018 on the basis of report from the international agency for research on cancer (IARC). Most common types of cancers occurring in humans are lung, female, breast and colorectal. There are high chances of death in these cancer patients on factors based on early detection chemo resistance or tumor heterogeneity as well as chances of metastases^[28]. Plants have been used as a source of effective anticancer agents, and 60% of currently used anticancer drugs are derived from natural sources such as plants, marine organisms, and microorganisms^[17]. Several studies have been conducted on herbs that possess anticancer properties and these have been used as potent anti-cancer drugs^[17]. Recently, the antitumor activity of Cinnamon has been shown to inhibit the proliferation of several human cancer cell lines including breast, leukemia, ovarian, and lung tumor cells^[17]. It is well known that metastasis, one of the major causes of mortality in cancer, involves cell adhesion, invasion, and migrate. In one study, the effect of CE on the migration of SiHa (human cervical tumor) cells was examined. The result showed that cinnamon significantly reduced the migration of cancer cells, demonstrating its potential use as an anticancer drug in cervical cancer^[17]. CE, which contains a mixture of polyphenols together with cinnamaldehyde as the major bioactive component, plays a therapeutic role in cervical cancer cells by depolarization of the mitochondrial membrane potential, resulting in cellular apoptosis^[17]. In another study, CE was shown to inhibit the growth of hematologic tumor cells. Moreover, the treatment of melanoma cell lines with CE also induced a decrease in

Cox-2 and HIF-1 α expression in the tumor tissues that mediate the potent antitumor activity of cinnamon. Cox-2 and HIF-1 α are well known as master regulators in cancer progression to aggravate angiogenesis and metastasis. The use of herbal medicine is supported by scientific studies for potent anticancer drugs. However, the use of an herbal medicine as an anticancer agent requires extensive research and strict criteria for standardization, safety, quality control, toxicity, and clinical trials [18].

Angiogenesis inhibitor

Angiogenesis is the development of new blood vessels in a mechanism used by tumors to promote growth and metastasis. VEGF is one of the most critical and specific angiogenesis factors. Various approaches for the inhibition of VEGF are currently being reviewed and studied in clinical trials [5]. Currently available anti-VEGF agents have been shown to have serious side effects such as hypertension, bleeding, and gastrointestinal perforation. There has been a renewed interest in identifying natural anti-VEGF agents, given their advantage of being safe for human use. Additionally, consumption of a plant-based diet has been effective in the prevention of cancer development [5]. One study has shown that Cinnamon is a natural diet-derived source of anti-VEGF agents [5]. It was found that cinnamon and its components (such as type A procyanidin trimer and tetrameric procyanidins) effectively inhibited the activity of VEGFR2 kinase as well as VEGF signaling in endothelial cells. The various aspects of angiogenesis, including endothelial cell proliferation, migration, as well as tumor-induced blood vessel formation in mice, were inhibited by CE. Further research is required to determine the major components responsible for the anti-angiogenic property of CE. As a natural inhibitor of VEGFR2, CE has the potential to be a routine diet-based strategy for cancer prevention and treatment [5].

Alzheimer's disease

Alzheimer's disease (AD) is a progressive, neurodegenerative, and irreversible brain disorder with symptoms such as memory loss, confusion, impaired judgments, and loss of language skills [19]. Many studies have proven that the accumulation of soluble oligomeric assemblies of β -amyloid polypeptides [amyloid-beta (Ab)] play a key role in AD development [19]. Compounds made from natural sources have been reported in many studies to inhibit the formation of Ab plaques [19]. A study found that Cinnamon extract (CEppt) inhibits the formation of toxic Ab oligomers and prevents the toxicity of Ab on neuronal PC12 cells. In another study, the oral administration of CEppt to an aggressive AD transgenic mice model led to the reduction of plaques and improvement in cognitive behavior. The results showed that the use of natural compounds such as cinnamon can inhibit toxic oligomeric Ab species formation in AD [19]. Extracellular plaques (such as Ab) and also intracellular neurofibrillary tangles of tau can cause the emergence of AD. Tangles are formed in later stages in relation to amyloid formation and are effective in the process of neurodegenerative progression [9]. Most of the drugs used for the prevention or treatment of AD are aimed to inhibit cholinesterase function or amyloid plaque formation. The agents capable of preventing the tau aggregation may be a key factor in the development of new drugs [9]. Tau is a protein that displays little to no structure

in regular conditions, and the molecule contains significant polyrich regions. The effects of an aqueous extract of cinnamon containing proanthocyanidins was examined on tau aggregation, and it was found that the extract of the whole cinnamon effectively inhibited the aggregation of human tau *in vitro*, and this could be attributed to both proanthocyanidin trimer and cinnamaldehyde in CE [9].

Antigastric ulcer

Helicobacter pylori is a major cause of chronic gastritis, peptic ulceration, and gastric cancer. Probiotic bacteria have been used for the inhibition of *H. pylori* and intestinal pathogens, prevention of inflammatory bowel diseases, reduction of the risks associated with mutagenicity and carcinogenicity, and improvement of the immune system [20]. Yogurt containing bifidobacterium and *Lactobacillus acidophilus* can suppress the growth of *H. pylori* in infected clinical patients [20]. Behrad *et al.* [20] showed that cinnamon yogurt exerted the strongest inhibitory effect on *H. pylori* growth *in vitro* in comparison with licorice yogurt and control yogurt for both strains tested [20]. The study showed that cinnamon preparations traditionally used for the treatment of gastrointestinal disorders are effective on suppressing *H. pylori* with little or no adverse reactions. These preparations could provide a potent anti-*H. pylori* effect that overcomes the problem of resistance to antibiotic treatments [21]. Simethicone is the medication most commonly used in treating dyspepsia. A study showed that the use of a cinnamon stomachic mixture is effective in alleviating the symptoms of dyspepsia by 70% and is similar to simethicone. The longer use of cinnamon stomachic mixture can increase the favorable response up to 80%, and it is considered safe, effective, and affordable compared to simethicone [22].

Antibacterial, antifungal

Cinnamon oil has the most potential bactericidal properties [16]. The antibacterial activity of cinnamon oil has been demonstrated, and it has been shown that cinnamon oil alone or in combination with triclosan, gentamicin, or chlorhexidine can effectively inhibit biofilm formation, detach existing biofilms, and kill bacteria in biofilms of clinical *Staphylococcus epidermidis* strains. Another advantage of the essential oil over antibiotics may be that bacteria do not develop resistance to essential oils.²³ However, further study is needed to clarify the complex mode of action of cinnamon and its components against biofilms of *S. epidermidis* and other clinically relevant microbes [23]. Cinnamaldehyde is one of the main components of cinnamon oil [23]. Cinnamaldehyde has been shown to destroy the cytoplasmic membrane of both Gram-positive and Gram-negative bacteria and induce depletion of the intracellular ATP concentration. Cinnamaldehyde inhibits the growth of fungi, including yeasts, filamentous molds, and dermatophytes, as well as the eggs and adults of human head lice, *Pediculus humanus capitis*. Aqueous and alcohol extracts of cinnamon also have demonstrated antibacterial effects against *H. pylori* [15].

Cinnamon osmophloeum Kaneh (Lauraceae), named as "indigenous cinnamon tree," is one of the hardwood species that possesses significant antifungal activity. *C. osmophloeum* is an endemic tree that grows in Taiwan's natural hardwood forest, and the chemical constituents of the essential oils found in its leaves are similar to those of *C.*

cassia bark oil. The leaf essential oils of *C. osmophloeum* were found to have strong inhibitory effects on the growth of fungi [24]. Cinnamaldehyde, the major compound in leaf essential oils, had the strongest antifungal activities compared with the other components. People can simply harvest the leaves to obtain a large amount of natural bioactive essential oils or cinnamaldehyde without debarking the trees; this could potentially lead to the development of wood decay preservation agents or fumigants in the future [24].

Nematicidal

There is a growing public concern for the environmental effects of pesticides on human health and the increase of undesirable effects on nontarget organisms by repeated applications of chemical pesticides [25]. Natural alternative sources such as plant essential oils have been suggested because they degrade to nontoxic products and have few harmful effects on nontarget organisms and the environment [25]. Pine wood nematode, *Bursaphelenchus xylophilus*, is associated with dead or dying conifers such as pines throughout North America and Japan [25]. In one study, the nematicidal activities of two cassia oils and four Cinnamon (肉桂 *ròu guì*) oils were evaluated, and the results showed that they are active against adult *B. xylophilus* and might be a good choice as a natural control agent [25]. The study showed that cinnamaldehyde is the most toxic compound against adult *B. xylophilus*, followed by (E)-2 methoxy cinnamaldehyde and (E)-cinnamic acid [25].

Acaricidal and repellent effect

House dust mites (*Dematophagoides farinae*, *Dematophagoides pteronyssinus*) are a major source of allergy in human households causing symptoms such as atopic dermatitis, asthma, perennial rhinitis, and even infant death syndrome. The use of chemical methods or treatment with repellents such as DEET and benzyl benzoate are related to the control of mite species [26]. However, the repeated use of these chemicals has resulted in the development of resistance against such chemicals and raised concerns regarding their effect on human health [26]. It is more effective to repel than to kill house mites because the allergen does not come from living dust mites but from their dead bodies or fecal particles [26]. To control house dust mites, it is better to find natural repellent materials such as plants with a range of bioactive chemicals [26]. The concentrated essential oil of Cinnamon has been tested for its acaricidal and repellent effect against house dust mites and has been shown to be a very effective material for this particular purpose, and could also reduce chemical resistance and harmful effects on human health [26].

Safety

Cinnamon has been used in food preparation and as medicinal remedies from ancient times. It is the most frequently consumed spice and is both safe and relatively inexpensive [14]. In one study, the intake of up to 6 g/d of *C. cassia* for > 40 days did not show any adverse effects. The most common adverse effects related to the excessive use of cinnamon were irritation and allergic reaction in skin or mucus membranes [15]. According to the United States Food and Drug Administration, *Cinnamon* spp., including common cinnamon and cassia cinnamon, are generally safe and well tolerated in amounts commonly found in food [15].

Cinnamon oil has also being listed and recognized as safe and is exempt from toxicity data requirements by the US Environmental Protection Agency [25].

Conclusion

Cinnamon has been utilised in many civilizations around the world as a spice for flavouring foods and as a natural folk medicine. Cinnamon has many specific functional properties such as antioxidant, anti-inflammatory, antimicrobial, and acaricidal activities, in addition to its medicinal values in relieving and treating a variety of serious illnesses such as diabetes, hyperlipidemia, gastrointestinal disorders, Alzheimer's disease, and cancer, according to the findings of various studies. More research is needed to determine the efficacy of cinnamon's active ingredients and their therapeutic benefits in the prevention and treatment of such disorders.

References

1. Vinitha M, Ballal M. *In vitro* anticandidal activity of *Cinnamomum verum*. J Med Sci. 2008;8:425e428.
2. Shan B, Cai YZ, Brooks DJ, Corke H. Antibacterial properties and major bioactive component of cinnamon stick (*Cinnamomum burmannii*): activity against food borne pathogenic bacteria. J Agric Food Chem 2007;55:5484e5490.
3. Ranasinghe P, Perera S, Gunatilake M, et al. Effects of *Cinnamomum zeylanicum* (Ceylon cinnamon) on blood glucose and lipids in a diabetic and healthy rat model. Pharmacognosy Res 2012;4:73e79.
4. Elshafie MM, Nawar IA, Algamal MA, Mohammad AS. Evaluation of the biological effects for adding cinnamon volatile oil and TBHQ as antioxidant on rats' lipid profiles. Asian J Plant Sci 2012;11:100e108.
5. Lu J, Zhang K, Nam S, Anderson RA, Jove R, Wen W. Novel angiogenesis inhibitory activity in cinnamon extract blocks VEGFR2 kinase and downstream signaling. Carcinogenesis 2010;31:481e488.
6. Molania T, Moghadamnia A, Aghel S, et al. The effect of cinnamaldehyde on mucositis and salivary antioxidant capacity in gamma-irradiated rats (a preliminary study). J Pharm Sci 2012;20:89.
7. Hong JW, Yang GE, Kim YB, Eom SH, Lew JH, Kang H. Anti-inflammatory activity of cinnamon water extract *in vivo* and *in vitro* LPS-induced models. BMC Complement Altern Med 2012;12:237.
8. Elahi RK. The effect of the cinnamon on dog's heart performance by focus on Korotkoff sounds. J Anim Vet Adv 2012;11:3604e3608.
9. Peterson DW, George RC, Scaramozzino F et al. Cinnamon extract inhibits tau aggregation associated with Alzheimer's disease *in vitro*. J Alzheimers Dis. 2009;17:585e597.
10. Safdar M, Khan A, Ali Khan Khattak MM, Siddique M. Effect of various doses of cinnamon on blood glucose in diabetic individuals. Pak J Nutr 2004;3:268e272.
11. Baker WL, Gutierrez-Williams G, White CM, Kluger J, Coleman C. Effect of cinnamon on glucose control and lipid parameters. Diabetes Care 2008;31:41e43.
12. Roussel A, Hininger I, Benaraba R, Ziegenfuss TN, Anderson RA. Antioxidant effects of a cinnamon extract in people with impaired fasting glucose that are overweight or obese. J Am Coll Nutr 2009;28:16e21.

13. Ziegenfuss TN, Hofheins JE, Mendel RW, Landis J, Anderson RA. Effect of a water- soluble cinnamon extract on body composition and features of the metabolic syndrome in pre-diabetic men and women. *J Int Soc Sports Nutr* 2006;3:45e53.
14. Anderson RA, Broadhurst C, Polansky MM, *et al.* Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin- like biological activity. *J Agric Food Chem* 2004;52:65e70.
15. Anubhav Dubey, Deepanshi Tiwari, Kshama Srivastava, Om Prakash, Rohit Kushwaha. A discussion on vinca plant. *J Pharmacogn Phytochem* 2020;9(5):27-31.
16. Prabuseenivasan S, Jayakumar M, Ignacimuthu S. *In vitro* antibacterial activity of some plant essential oils. *BMC Complement Altern Med* 2006;6:39.
17. Koppikar SJ, Choudhari AS, Suryavanshi SA, Kumari S, Chattopadhyay S, Kaul- Ghanekar R. Aqueous cinnamon extract (ACE-c) from the bark of *Cinnamomum Cassia* causes apoptosis in human cervical cancer cell line (SiHa) through loss of mitochondrial membrane potential. *BMC Cancer*. 2010;10:210.
18. Kwon H, Jeon WK, Hwang J, *et al.* Cinnamon extract suppresses tumor progression by modulating angiogenesis and the effector function of CD8^b T cells. *Cancer Lett* 2009;278:174e182.
19. Frydman-Marom A, Levin A, Farfara D, *et al.* orally administrated cinnamon extract reduces b-Amyloid oligomerization and corrects cognitive impairment in Alzheimer's disease animal models. *PLoS One*. 2011;28:e16564.
20. Behrad S, Yusof MY, Goh KL, Baba AS. Manipulation of probiotics fermentation of yogurt by cinnamon and licorice: effects on yogurt formation and inhibition of *Helicobacter pylori* growth *in vitro*. *World Acad Sci Eng Technol*. 2009;60:590e594.
21. Ahmed MA. Efficacy of ginger and cinnamon pharmaceutical preparations in patients with *Helicobacter pylori*-associated functional dyspepsia. *Prime J Microbiol Res* 2012;2:67e72.
22. Jindarat S, Muangnoi C, Changsiriporn D, *et al.* Efficacy and safety of cinnamon stomachic mixture for patients with functional dyspepsia. *Siriraj Med J*. 2006;58:1103e1106.
23. Nuryastuti T, Van Der Mei HC, Busscher HJ, Irvati S, Aman AT, Korm BP. Effect of cinnamon oil on IcaA expression and biofilm formation by *Staphylococcus epidermidis*. *Appl Environ Microbiol*. 2009;75:6850e6855.
24. Wang S, Chen P, Chang S. Antifungal activities of essential oils and their constituents from indigenous cinnamon (*Cinnamomum osmophloeum*) leaves against wood decay fungi. *Bioresour Technol*. 2005;96:813e818.
25. Kong JO, Lee SM, Moon YS, Ahn YJ. Nematicidal activity of cassia and cinnamon oil compounds and related compounds toward *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae). *J Nematol*. 2007;39:31e36.
26. Oh MJ. The acaricidal and repellent effect of cinnamon essential oil against house dust mite. *World Acad Sci Eng Technol*. 2011;60:710e714.
27. Dugoua J, Seely D, Perri D, *et al.* From type 2 diabetes to antioxidant activity: a systematic review of the safety and efficacy of common and cassia cinnamon bark. *Can J Physiol Pharmacol* 2007;85:837e847.
28. AnubhavDubey, Deepanshi Tiwari, Yatendra Singh, Om Prakash, Pankaj Singh. Drug repurposing in Oncology: Opportunities and challenges.. *Int J of Allied Med Sci and Clin Res* 2021;9(1):68-87.
29. Dubey Anubhav, Tiwari Mamta, Kumar Vikas, Srivastava, Kshama, Singh, Akanksha. Investigation of anti-hyperlipidemic activity of vinpocetine in wistar rat *International Journal of Pharmaceutical Research* 2020;12(02):1879-1882.