E-ISSN: 2707-2835
P-ISSN: 2707-2827
www.pharmacognosyjournal.com
IJPLS 2021; 2(2): 01-05
Received: 01-03-2021
Accepted: 03-07-2021

Lisette H Degla
(1) Zootechnical Research and Livestock System Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin
(2) Laboratory of Biochemistry and Bioactive Natural Substances, Faculty of Science and Technology, University of Abomey-Calavi, 04BP 0320, Cotonou, Benin

Pascal A Olounlade
(1) Zootechnical Research and Livestock System Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin
(2) Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, 01 BP 526 Cotonou, Benin

Sabbas Attindehou
Animal Health and Biosecurity Research Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin

Mawulé S Hounzangbe-Adote
Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, 01 BP 526 Cotonou, Benin

Latifou Lagnika
Laboratory of Biochemistry and Bioactive Natural Substances, Faculty of Science and Technology, University of Abomey-Calavi, 04BP 0320, Cotonou, Benin

Corresponding Author:
Pascal Abiodoun OLOUNLADE
(1) Zootechnical Research and Livestock System Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin
(2) Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, 01 BP 526 Cotonou, Benin

Comparative study of acute oral toxicity of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* in wistar rats

Lisette H Degla, Pascal A Olounlade, Sabbas Attindehou, Mawulé S Hounzangbe-Adote and Latifou Lagnika

DOI: <https://doi.org/10.33545/27072827.2021.v2.i2a.32>

Abstract

The essential oils from *Ocimum gratissimum*, *Hyptis suaveolens* (Lamiaceae) and *Psidium guajava* (Myrtaceae) have biological properties. The aim of this work is to compare the effect of essential oils of these three medicinal plants on weight growth, hematological and biochemical parameters of treated rats through the procedure of acute toxicity evaluation on Wistar rats. The doses applied are 100, 250 and 500mg/Kg bw for each of the essential oils. From the results obtained, the effects of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* leaves on biochemical, hematological parameters and weight growth of rats are dose dependent. At the dose of 500mg/Kg bw, they significantly increased the rate of white blood cells, urea, creatinemia, transaminases and reduced the weight growth of rats by more than 10% ($p < 0.05$) compared to the control lot. This shows that these essential oils can have toxic effects when applied in high doses.

Keywords: Essential oils, Biochemical and Hematological parameters, Weight growth, Toxicity

Introduction

The use of medicinal plants in the treatment of diseases is a practice that has existed since ancient times [23]. A great importance is more and more given to this science because of the health problems that remain unanswered or that have difficulties, in particular, the resistance of pathogenic microorganisms to conventional drugs [24]. Many of these medicinal plants have proven biological properties. These scientific researches will have reached their goals when the population would have found relief and satisfaction through the use of medicinal plants, or its derivatives (Extracts, Essential Oils), and or TAMs (Traditionally Improved Medicines). Thus, it is important to conduct safety studies of plant species with biological properties to ensure that their uses will not have a negative impact on health. To this end, this work proposes to compare the toxicity of essential oils of three medicinal plants *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava*. *Psidium guajava* is a shrub of the Myrtaceae family whose fruits are edible and whose leaves are used in therapy. *Ocimum gratissimum* is an herb of the Lamiaceae family increasingly cultivated in gardens for its nutritional and therapeutic values. Found at the edge of the roads in the villages and qualified as weeds, *Hyptis suaveolens* of the Lamiaceae family is also solicited in traditional medicine. In sum, *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* are used in the treatment of dysentery, diarrhea vomiting, rheumatism, diabetes, and gastroenteritis, pulmonary problems, to cure wounds, ulcer, rheumatism, fever, malaria and intestinal parasitosis [1, 2]. They are used in the form of decoction, infusion or macerated but the dosage is often unclear. The method of preparation (decoction, maceration, infusion), the type of solvent used (water, ethanol, sugar beverages, carbonated water) can extract one or the other of the secondary metabolites responsible for the biological activities. However, essential oils, which are also secondary metabolites of medicinal plants, escape these methods and techniques of extraction. And yet, they possess interesting biological properties [25, 26] and act at very low quantities. It has been reported that essential oils extracted from the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* have Antioxidant, antibacterial, antifungal, insecticidal, anti-inflammatory, antidiarrheal and antiparasitic properties [3-8].

They can be used in the treatment of infectious diseases, as bioinsecticide and food industries. Essential oils are composed of several molecules that can act in synergy or the biological activity can only depend on the majority molecules [22, 26]. It is with regard to all these observations that the present study proposes to study the safety of essential oils of the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava*.

Material

Fresh leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* are collected in December 2020 in Abomey-calavi (Benin). Herbariums of each of the plants were authenticated at the National Herbarium of the University of Abomey-calavi and registered under the numbers YH 603/HNB, YH 604/HNB and YH 605/HNB respectively for *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava*. 30 healthy, non-pregnant, nulliparous female rats weighing between 150 and 200g and 8 to 12 weeks of age are used for acute toxicity tests. They were obtained at the animal house of the National University of Agriculture of Kétou and are kept in a room with artificial lighting with a sequence of 12 hours of light and 12 hours of darkness with a temperature between 23 °C and 25 °C and a relative humidity of 40 to 45. They were fed with Veto- service® pellets and had unlimited access to tap water.

Method

Preparation of essential oil extracts

The essential oils of the leaves are obtained through the extraction of a known mass of fresh leaves of each species by hydrodistillation for 3h using a Clevenger type device. The essential oils collected in vials are stored at 4 °C. For the evaluation of the tests, the essential oils were prepared in a hydroalcoholic solution.

Acute toxicity

The determination of acute toxicity was conducted according to OECD guideline 423₍₂₀₎. The rats were divided into 10 groups of 03 each. The Control lot received 1 mL of the hydroalcoholic solution. Lots 1, 2 and 3 received the essential oil of *Ocimum gratissimum* leaves at doses of 100, 250 and 500 mg/Kg bw respectively. The essential oil of *Hyptis suaveolens* leaves was administered to the rats of lots 4, 5 and 6 respectively at the doses of 100, 250 and 500

mg/Kg bw. And 100, 250 and 500 mg/Kg bw of *Psidium guajava* essential oil was given to lots 7, 8 and 9 respectively. Before the tests, the rats were deprived of food but not water overnight. Then they were weighed and the essential oils were administered through a stomach tube. The rats were fed and put under observation for 14 days. Weight monitoring was performed at D0, D7 and D14. At the end of the 14 days, blood samples were taken for hematological and biochemical analysis.

Statistical Analysis

To assess the effect of essential oils and dose on weight, biochemical and hematological parameters, an analysis of variance was performed with R 4.0.3 softwares (R Core Team, 2020) in case of normality. When normality was not acquired, the nonparametric Kruskal Wallis test was performed. Mean structuring was done with the SNK function of the "agricolae" package [21] when the probability was significant ($p < 0.05$).

Results

No signs of toxicity such as agitation, convulsions, drowsiness and or diarrhea were observed in the different lots of rats treated with the essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* leaves. This indicates that the essential oils of the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* have an LD50 greater than 500mg/kg bw.

Effect of essential oils on the growth of treated rats

Table 1, 2 and 3 show the effect of administration of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens*, and *Psidium guajava* respectively on the weight growth of rats. Treatment with the essential oil of *Ocimum gratissimum* leaves significantly reduced the growth rate of rats at the dose of 500mg/Kg bw after 14 days ($p < 0.05$). Treatment with the essential oil of *Psidium guajava* leaves is associated with a significant reduction in the growth rate of rats ($p < 0.05$). The essential oil of *Hyptis suaveolens* leaves stimulated the growth of rats between Day 1 and Day 7. However, the best growth rate is obtained with the dose of 250mg/Kg bw at the end of 14 days although in the interval of days D7-D14, there is no increase in the weight mass of rats.

Table 1: Effect of essential oil of *Ocimum gratissimum* leaves on weight growth of treated Wistar rats.

EO	Dose (mg/Kg bw)	Weight growth (g)		
		D1-D7	D1-D14	D7-D14
OG	500	0.03 ^b	0.03 ^b	0 ^b
	250	0.24 ^a ±0.02	0.29 ^a ±0.02	0.04 ^c ±0.0003
	100	0.13 ^a ±0.0001	0.16 ^c ±0.01	0.03 ^c ±0.000
Control	-	0.19 ^a ±0.001	0.36 ^a ±0.002	0.13 ^a ±0.00003

a, b, c averages in the same row with different superscript as significantly different ($P < 0.05$). OG: *Ocimum gratissimum*; EO: essential oil; D1: Day 1; D7: Day 7; D14: Day 14

Table 2: Effect of essential oil of *Psidium guajava* leaves on weight growth of treated Wistar rats.

EO	Dose (mg/Kg bw)	Weight growth (g)		
		D1-D7	D1-D14	D7-D14
PG	500	0.21 ^a ±0.001	0.26 ^b ±0.006	0.04 ^b ±0.0005
	250	0.11 ^b ±0.001	0.20 ^b ±0.004	0.08 ^b ±0.0005
	100	0.08 ^c ±0.001	0.14 ^c ±0.0003	0.06 ^b
Control	-	0.19 ^a ±0.001	0.36 ^a ±0.002	0.13 ^a ±0.00003

a, b, c averages in the same row with different superscript as significantly different ($P < 0.05$). PG: *Psidium guajava*; EO: essential oil; D1: Day 1; D7: Day 7; D14: Day 14

Table 3: Effect of essential oil of *Hyptis suaveolens* leaves on weight growth of treated Wistar rats.

EO	Dose (mg/Kg bw)	Weight growth (g)		
		D1-D7	D1-D14	D7-D14
HS	500	0.35 ^b ±0.0008	0.37 ^a ±0.006	0.02 ^b ±0.002
	250	0.47 ^b ±0.03	0.45 ^a ±0.02	-0.01 ^b ±0.0028
	100	0.26 ^a ±0.0007	0.40 ^a ±0.005	0.11 ^a ±0.002
Control	-	0.19 ^a ±0.001	0.36 ^a ±0.002	0.13 ^a ±0.00003

a, b, c averages in the same row with different superscript as significantly different ($P<0.05$). D1: Day 1; D7: Day 7; D14: Day 14

Effect of essential oils on hematological and biochemical parameters of treated Wistar rats

According to the analysis of hematological and biochemical parameters, the administration of the essential oil of the leaves of *Ocimum gratissimum*, is associated with an increase in the level of blood white blood cells, creatinemia, urea and transaminases compared to the control lot (Table 4). The effect of the essential oil on these parameters is dose dependent ($p<0.05$). Similarly, administration of the essential oil of *Hyptis suaveolens* leaves was associated with

a significant difference ($p<0.05$) in white blood cell and urea levels compared to the control lot (Table 5). At doses 250 and 500mg/Kg bw, the essential oil of *Psidium guajava* leaves induced elevated white blood cell and urea levels in the treated rat lots compared to the control lot (Table 6). The elevated values of transaminases (AST and ALT) are obtained in rats treated with 500mg/Kg bw of the essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* leaves.

Table 4: Effect of essential oil of *Ocimum gratissimum* leaves on hematological and biochemical parameters of Wistar rats

OG (mg/Kg bw)	Urea (g/L)	Creat (mg/L)	AST (IU)	ALT (UI)	WBC ($\times 10^9/L$)	RBC ($\times 10^6/mm^3$)	Ht (%)	Hb (g/dL)
100	8.77 ^b ±0.18	0.41 ^b ±0.04	18.63 ^d ±1.61	161.8 ^c ±55.72	4.52 ^c ±0.10	7.40 ^b ±0.17	47.75 ^a ±1.13	14.55 ^a ±0.43
250	7.87 ^b ±0.54	0.36 ^b ±0.01	21.83 ^c ±0.51	221.42 ^b ±171.47	5.48 ^b ±0.17	7.42 ^b ±0.05	45.55 ^a ±1.01	14.25 ^a ±0.14
500	9.88 ^c ±0.01	0.43 ^b	50.63 ^b ±0.03	257.42 ^b ±8.80	5.85 ^b ±0.13	7.60 ^b ±0.23	45.00 ^a ±0.17	14.30 ^a
Control	5.85 ^a ±0.38	0.23 ^a ±0.03	55.00±28.70	116.10 ^a ±16.60	3.15 ^a ±0.26	5.25 ^a ±0.05	44.65 ^a ±0.43	15.30 ^a ±0.29

a, b, c, d averages in the same row with different superscript as significantly different ($P<0.05$). OG: *Ocimum gratissimum*, AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cell; RBC: Red Blood Cells; Ht: Hematocrit; Hb: Hemoglobin.

Table 5: Effect of essential oil of *Hyptis suaveolens* leaves on hematological and biochemical parameters of Wistar rats

HS (mg/Kg/bw)	Urea (g/L)	Creat (mg/L)	AST (IU)	ALT (UI)	WBC ($\times 10^9/L$)	RBC($\times 10^6/mm^3$)	Ht (%)	Hb (g/dL)
100	4.49 ^c ±0.14	0.22 ^a ±0.05	28.81 ^c ±8.57	64.92 ^d ±5.23	4.40 ^c ±0.12	4.59 ^a ±0.34	39.00 ^a ±2.89	14.25 ^a ±0.84
250	5.75 ^a ±0.11	0.25 ^a ±0.03	67.80 ^b ±35.50	220.00 ^b ±26.20	5.20 ^b ±0.07	5.33 ^a ±0.10	43.90 ^a ±0.06	15.00 ^a ±0.06
500	8.67 ^b ±0.12	0.20 ^a ±0.01	75.10 ^b ±12.10	300.30 ^c ±65.80	6.55 ^d ±0.78	7.60 ^b ±0.46	44.25 ^a ±0.61	15.55 ^a ±0.43
Control	5.85 ^a ±0.38	0.23 ^a ±0.03	55.00±28.70	116.10 ^a ±16.60	3.15 ^a ±0.26	5.25 ^a ±0.05	44.65 ^a ±0.43	15.30 ^a ±0.29

a, b, c averages in the same row with different superscript as significantly different ($P<0.05$). HS: *Hyptis suaveolens*, AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cell; RBC: Red Blood Cells; Ht: Hematocrit; Hb: Hemoglobin.

Table 6: Effect of essential oil of *Psidium guajava* leaves on hematological and biochemical parameters of Wistar rats

PG (mg/Kg/bw)	Urea (g/L)	Creat (mg/L)	AST (IU)	ALT (UI)	WBC ($\times 10^9/L$)	RBC($\times 10^6/mm^3$)	Ht (%)	Hb (g/dL)
100	5.46 ^a ±0.89	0.24 ^a ±0.01	29.5 ^c ±9.3	50.60 ^c ±16.10	2.25 ^d ±0.03	5.11 ^a ±0.06	44.15 ^a ±0.09	14.10±0.06
250	6.00 ^a ±0.16	0.19 ^c ±0.02	39.21 ^a ±0.63	131.00 ^a ±64.50	4.60 ^c ±0.52	5.12 ^a ±0.05	43.55 ^a ±0.43	14.30±0.06
500	10.65 ^b ±0.21	0.48 ^b ±0.02	48.18 ^b ±10.11	209.10 ^b ±80.50	5.20 ^b ±0.56	8.17 ^b ±0.11	50.15 ^b ±1.47	15.00±0.52
Control	5.85 ^a ±0.38	0.23 ^a ±0.03	55.00±28.70	116.10 ^a ±16.60	3.15 ^a ±0.26	5.25 ^a ±0.05	44.65 ^a ±0.43	15.30 ^a ±0.29

a, b, c, d averages in the same row with different superscript as significantly different ($P<0.05$). PG: *Psidium guajava*, AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cell; RBC: Red Blood Cells; Ht: Hematocrit; Hb: Hemoglobin.

Comparison of the effects of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* on hematological and biochemical parameters of treated rats

Table 7 compares the effect of administration of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* with each other on the renal parameters of treated lots of wistar rats. There was no significant

difference between OG, HS and PG on transaminases and hemoglobin levels of treated batches of rats ($p>0.05$). However, the OG treatment increased urea, creatinemia and red blood cell levels more. PG, and HS treatments showed similar activity on blood urea and red blood cell levels. The OG and HS treatments showed similar activity on blood white blood cell count.

Table 7: Comparison of the effects of essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* on hematological and biochemical parameters of treated rats

Parameters	Essential oils				p-value
	Témoïn	HS	OG	PG	
Urea (mg/L)	5.85 ^b ±0.38	6.30 ^{ab} ±0.62	8.84 ^a ±0.33	7.37 ^{ab} ±0.87	0.030
Creatinemia (mg/L)	0.23 ^b ±0.03	0.22 ^b ±0.02	0.40 ^a ±0.02	0.30 ^{ab} ±0.05	0.002
AST (UI)	116.10±16.60	195.10±44.90	141.98±7.85	209.10±80.50	0.111
ALT (UI)	55.00±28.70	57.20±13.20	53.00±16.10	65.20±25.90	0.973
WBC(x10 ⁹ /L)	3.15 ^b ±0.26	5.38 ^a ±0.39	5.28 ^a ±0.21	4.02 ^{ab} ±0.50	0.007
RBC(x10 ⁹ /mm ³)	5.25 ^b ±0.05	5.84 ^{ab} ±0.48	7.53 ^a ±0.10	6.13 ^{ab} ±0.51	0.012
HB (g/dl)	15.30±0.29	14.93±0.33	14.37±0.14	14.47±0.20	0.133
Ht (%)	44.65±0.43	42.38±1.20	46.10±0.61	45.95±1.14	0.050

a, b, c, d averages in the same row with different superscript as significantly different ($P<0.05$). OG: *Ocimum gratissimum*; HS: *Hyptis suaveolens*; PS: *Psidium guajava*; AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cells; RBC: Red Blood Cells; Ht: Hematocrit; Hb: Hemoglobin.

Discussion

Toxicity studies make it possible to evaluate the short, medium and long term impacts that the intake of a given substance could have on the organism. In the control of gastrointestinal parasitosis of small ruminants and humans, the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* (three aromatic medicinal plants) are used. They could be an alternative in the fight against parasite resistance to synthetic anthelmintics and reduce ecotoxicity. In the present study, the acute toxicity of the essential oils of these three medicinal plants were evaluated on Wistar rats at different doses (100; 250 and 500 mg/Kg bw) in order to evaluate the impacts that their uses could have on the organism. According to the work of Bezerra [9], on *Artemia salina* larvae, the essential oil of *Hyptis suaveolens* leaves can be toxic. Similar results had been reported instead by Kpadonou [10] for the essential oil of *Ocimum gratissimum*. According to the Work of Fandohan [11], the LD50 of *Ocimum gratissimum* essential oil is 1750 mg/Kg bw. However, only the rats having received doses of 5-500 mg/Kg bw not shown any sign of toxicity like torpid and death. This justifies the choice of the doses used in this study.

In the present study, in general, it was observed that the essential oils of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* leaves influenced the growth of rats, their hematological parameters (White blood cell count) and biochemical parameters (Urea, creatinemia, transaminases) after 14 days of observations compared to the untreated lot. On the growth of rats, the essential oil of *Hyptis suaveolens* leaves stimulated a weight gain of the animals but with a low rate at the dose of 500mg/Kg bw. The essential oils of *Ocimum gratissimum* and *Psidium guajava* leaves, compared to the control lot, significantly ($p<0.05$) decreased the growth rate of the rats with more than 10% reduction at the dose of 500mg/Kg bw. These results suggest that the essential oils of the leaves of *Psidium guajava* and *Ocimum gratissimum*, can be potentially toxic. Indeed, a significant weight loss of more than 10% is a sign of toxic effects caused by the intake of a substance [12, 13]. It could be explained by the presence in these essential oils of anti-nutritional compounds responsible for a decrease in food intake of treated rats [14]. According to the work of Efosa and Udoha [15, 16], repeated high dose administration of *Ocimum gratissimum* leaf extract, can be toxic. However, it followed from the work of [12, 17], that aqueous and methanolic extracts of *Psidium guajava* leaves are free from toxicity. However, from the dose 1200 mg/Kg bw, some signs of toxicity of the ethanolic extract were observed by Manikandan [18] who estimated 600mg/Kg bw as the non-toxic dose. The essential oil of *Ocimum gratissimum* leaves, acted on the hematopoietic system of rats by a

significant ($p<0.05$) increase in the blood white blood cell count. These results corroborate [16]. The same phenomenon was induced by the essential oil of *Psidium guajava* and *Hyptis suaveolens* leaves at the dose of 500 mg/Kg bw. The increase in the number of blood white blood cells is observed in inflammatory processes [14, 19]. Urea an important nitrogenous metabolic waste product released during protein metabolism and creatinine produced and released in body fluids showed an increase in their levels in lots of rats treated with the essential oils at doses of 250 and 500mg/Kg bw. They are biomarkers for determining renal function. An increase in their concentration in the blood indicates an alteration of the renal function and also the toxic effects that a compound could have [18]. An increase in transaminase levels is observed at the dose of 500mg/Kg bw may suggest liver damage. Essential oils from the leaves of *Ocimum gratissimum*, *Psidium guajava* and *Hyptis suaveolens* may have beneficial biological effects on organisms however at high doses and this from 500mg/Kg bw they showed signs of toxicity. The high toxicity of essential oils can be explained by their lipophilic character which makes them capable of damaging the plasma membrane by affecting the structure of polysaccharides, the phospholipids [9]. The complexity of the chemical composition of the essential oils of the leaves of these three medicinal plants can explain the different results obtained. However, some factors such as metabolism and bioavailability of these essential oils which are not taken into account in the present work are important to explain the mechanisms of action of these essential oils in the organism of animals and human.

Conclusion

In sum, the essential oils of the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava* may show signs of toxicity above a level. Any abusive use should be avoided. Essential oils give satisfactory results at very small doses compared to the doses used in this study. The complexity of the chemical composition of the essential oils of medicinal plants and their mechanisms of action may explain the variations obtained in the results obtained from lots of rats treated with the essential oils of the leaves of *Ocimum gratissimum*, *Hyptis suaveolens* and *Psidium guajava*. Histological analyses of the organs and complementary studies of chronic and sub-chronic toxicity will be necessary for a good use of the essential oils of these medicinal plants.

References

1. Degla LH, Olounlade PA, Amoussa AMO, EVB Azando, Hounzangbe-Adote MS, *et al.* Pharmacognostical, biochemical activities and zootechnical applications of *Psidium guajava* (Myrtaceae), plant with high medicinal value in tropical and subtropical parts of the World: A

- review. J. Med. Plants Stud. 2021a;9(3):14-18.
2. Degla LH, Olounlade PA, Amoussa AMO, EVB Azando, Hounzangbe-Adote MS *et al.* Pharmacological and Biochemical Aspects of the Lamiaceae Family used in the Treatment of Intestinal Parasitosis in West and Central Africa. Pharmacogn Rev 2021b ;15(29):69-75.
 3. Arain A, Sherazi STH, Mahesar SA, Sirajuddin. Essential Oil From *Psidium guajava* Leaves: An Excellent Source of β -Caryophyllene. Nat. Prod. Commun. 2019;14 :1-5.
 4. Ngbolua K, Lufuluabo LG, Moke LE, Bongo GN, Liyongo CI, *et al.* A review on the Phytochemistry and Pharmacology of *Psidium guajava* L. (Myrtaceae) and Future direction. Discov. Phytomed 2018;5(2):7-13.
 5. Tulugu M, Sujatha B, Lakshmi BS. Anatomical, phytochemical and *in vitro* antimicrobial studies of *Hyptis suaveolens* L. Poit. of family Lamiaceae. Int J Bio-Pharm Res 2019;8(6):2617-22.
 6. Castro DJAM, Monteiro OS, Coutinho DF, Rodrigues AAC, Silva DJKR, *et al.* Seasonal and Circadian Study of a Thymol/ γ -Terpinene/p-Cymene Type Oil of *Ocimum gratissimum* L. and Its Antioxidant and Antifungal Effect. J Braz Chem Soc 2019;30(5):930-8.
 7. Olukowade IL, Elusiyan CA, Omisore NO, Adewoyin FB, Adewunmi CO. A comparative study of the anti-protozoan activities of *Hyptis suaveolens* (L.) Poit and *Hyptis atrorubens* (L.) Poit. World Sci. News 2021;159:71-80.
 8. Machado FDF, Formiga RO, Lima GRM, de Jesus NZT, Júnior EBA, *et al.* *Hyptis suaveolens* (L.) Poit protects colon from TNBS-induced inflammation via immunomodulatory, antioxidant and anti-proliferative mechanisms. J. Ethnopharmacol 2021;265 :113153, 1-13.
 9. Bezerra JWA, Costa AR, da Silva MAP, Rocha MI, Boligon AA, *et al.* Chemical composition and toxicological evaluation of *Hyptis suaveolens* (L.) Poiteau (Lamiaceae) in *Drosophila melanogaster* and *Artemia salina*. S. Afr. J. of Bot 2017 ;113:437-442.
 10. Kpadonou-Kpoviessi BGH, Kpoviessi DSS, Yayi-Ladekan E, Gbaguidi F, Yehouenou B, *et al.* Phytochemical screening, antimicrobial activities and toxicity against *Artemia salina* Leach of extracts and fractions of *Ocimum gratissimum* Linn from Benin. J. Chem. Pharm. Res 2013;5(10):369-376.
 11. Fandohan P, Gnonlonfin B, Laleye A, Gbenou JD, Darboux R, *et al.* Toxicity and gastric tolerance of essential oils from *Cymbopogon citratus*, *Ocimum gratissimum* and *Ocimum basilicum* in Wistar rats. Food Chem. Toxicol 2008;46:2493-2497.
 12. Babatola LJ, Oboh G, Ademiluyi AO. Toxicological evaluation of aqueous extract of different varieties of guava (*Psidium guajava* Linn) leaves. Comp. Clin. Pathol 2019. Doi : 10.1007/s00580-019-03002-0.
 13. Atik N, Muda I, Rahmadi AR, Achadiyani, Djunaedy DD. Phytochemical screening and histology appearance of acute oral toxicity study on ethanol extract of *Psidium guajava* Linn. Fruit in mice. Asian J Pharm Clin Res 2019;12(1):351-355.
 14. Manekeng HT, Mbaveng AT, Mendo SAN, Agokeng AJD, Kuete V. Evaluation of Acute and Subacute Toxicities of *Psidium guajava* Methanolic Bark Extract: A Botanical with *in vitro* Antiproliferative Potential. Evid. Based Complement. and Alternat. Med 2019, 1- 13.
 15. Efosa OB, Osarogie OJ. Histo-morphological Examination of the Visceral Organs of Albino Wistar Rats pre-exposed to *Ocimum gratissimum* Crude Decoction. Int. J. Pharm. Sci. Invent 2014;3(5):36-41.
 16. Udoha MU, Okolie NJ, Ijioma SN. Chronic Oral Administration of *Ocimum gratissimum* Leaf Extract Induced Gastrointestinal Necrosis in Treated Albino Rats. EJMP 2019;27(3):1-9.
 17. Yamssi C, Payne VK, Noumedem Anangmo CN, Tateng Ngouateu A, Megwi L *et al.* Acute Toxicity of *Pentaclethra macrophylla* and *Psidium guajava* Use as Antiprotozoan Medicinal Plants. J Drug Discov Develop and Deliv 2020;6(1):1037.
 18. Manikandan R, Vijaya Anand A, Rengarajan RL, Velayuthaprabhu S, Bharathi V. Acute toxicity and dose fixation of the ethanolic leaf extract of *Psidium guajava*. J. Nat. Remedies 2020;4(S2). 21
 19. Olounladé PA, Konmy BBS, Azando EVB, Baba-Moussa L, Allou SD. Feeding Rabbits (*Oryctolagus cuniculus*) Based on *Ocimum gratissimum* and *Vernonia amygdalina* Leaves in West Africa: Biochemical and Hematological Aspects. Biochem. Mol. Biol. 2021;6(3)40-47. doi: 10.11648/j.bmb.20210603.12.
 20. OCDE 423. Ligne directrice de l'OCDE pour les essais de Produits chimiques 2001, 1-14.
 21. Mendiburu F. agricolae: Statistical Procedures for Agricultural Research. R package version 1.3-3 2020. <https://CRAN.R-project.org/package=agricolae>. Abidi A, Sebai E, Dhibi M, Alimi D, Rekik M, *et al.* Chemical analyses and anthelmintic effects of *Artemisia campestris* essential oil, Vet. Parasitol. 2019 ;<https://doi.org/10.1016/j.vetpar.2018.10.003>
 22. Abdulsalami H, Mudi SY, Daudu O, Abdulazeez Y, Aliyu BS, *et al.* Ethnobotanical survey of medicinal plants used in the treatment of gastrointestinal tract infections in Ebiraland Kogi state, Nigeria. J. Med. Plants Stud 2020;8(1):38-44.
 23. Dounon V, Hounsa E, Koudokpon H, Legba BB, Fabiyi K, *et al.* A Literature Review-*Khaya senegalensis*, *Anacardium ouest* L., *Cassia sieberiana* DC., *Pterocarpus erinaceus*, *Diospyros mespiliformis*, *Ocimum gratissimum*, *Manihot esculenta*, *Vernonia amygdalina* Delile, *Pseudocedrela kotschy* and *Daniellia oliveri* Possess Properties for Managing Infectious Diarrhea. ABB 2020;11:457-473. <https://doi.org/10.4236/abb.2020.1110031>.
 24. Tohidi B, Rahimmalek M, Arzani A. Essential oil composition, total phenolic and flavonoid contents, and antioxidant activity of Thymus species collected from different regions of Iran, Food Chem 2016; doi: <http://dx.doi.org/10.1016/j.foodchem.2016.09.20>.
 25. Pandey AK, Singh P, Tripathi NN. Chemistry and bioactivities of essential oils of some *Ocimum* species: an overview. Asian Pac J Trop Biomed 2014;4(9):682-694.